



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Travelers Can Import Colistin-Resistant Enterobacteriaceae, Including Those Possessing the Plasmid-Mediated *mcr-1* Gene

Bernasconi, Odette J ; Kuenzli, Esther ; Pires, João ; Tinguely, Regula ; Carattoli, Alessandra ; Hatz, Christoph ; Perreten, Vincent ; Endimiani, Andrea

DOI: <https://doi.org/10.1128/AAC.00731-16>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-128433>

Journal Article

Published Version

Originally published at:

Bernasconi, Odette J; Kuenzli, Esther; Pires, João; Tinguely, Regula; Carattoli, Alessandra; Hatz, Christoph; Perreten, Vincent; Endimiani, Andrea (2016). Travelers Can Import Colistin-Resistant Enterobacteriaceae, Including Those Possessing the Plasmid-Mediated *mcr-1* Gene. *Antimicrobial Agents and Chemotherapy*, 60(8):5080-5084.

DOI: <https://doi.org/10.1128/AAC.00731-16>

Travelers Can Import Colistin-Resistant *Enterobacteriaceae*, Including Those Possessing the Plasmid-Mediated *mcr-1* Gene

Odette J. Bernasconi,^{a,b} Esther Kuenzli,^{c,d} João Pires,^{a,b} Regula Tinguely,^a Alessandra Carattoli,^{a,e} Christoph Hatz,^{c,d}
Vincent Perreten,^f Andrea Endimiani^a

Institute for Infectious Diseases, University of Bern, Bern, Switzerland^a; Graduate School for Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland^b; Swiss Tropical and Public Health Institute, Basel, Switzerland^c; Division of Communicable Diseases, Institute for Social and Preventive Medicine, University of Zurich, Zurich, Switzerland^d; Department of Infectious, Parasitic and Immune-Mediated Diseases, Istituto Superiore di Sanità, Rome, Italy^e; Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Bern, Switzerland^f

Stool samples from 38 travelers returning from India were screened for extended-spectrum cephalosporin- and carbapenem-resistant *Enterobacteriaceae* implementing standard selective plates. Twenty-six (76.3%) people were colonized with CTX-M or DHA producers, but none of the strains was colistin resistant and/or *mcr-1* positive. Nevertheless, using overnight enrichment and CHROMagar Orientation plates supplemented with colistin, four people (10.5%) were found to be colonized with colistin-resistant *Escherichia coli*. One cephalosporin-susceptible sequence type 10 (ST10) strain carried a 4,211-bp IS*Apl1-mcr-1*-IS*Apl1* element in an IncHI2 plasmid backbone.

The emergence of colistin-resistant (COL-R-Ent) *Enterobacteriaceae* harboring the plasmid-mediated *mcr-1* gene has raised serious concerns (1, 2). These strains (especially *Escherichia coli*) have been isolated worldwide from humans (1, 3–10), food-producing animals (4, 6, 11–14), the food chain (7, 10, 15–17), and the environment (4, 15). However, most of these studies searched for *mcr-1* in previously stored extended-spectrum cephalosporin-resistant *Enterobacteriaceae* (ESC-R-Ent) (8, 12, 13, 15, 17). This leads to a question about the actual prevalence of *mcr-1*-positive *Enterobacteriaceae* (*mcr-1*-Ent), especially because *mcr-1* can be carried by plasmids not coharboring extended-spectrum β -lactamase (ESBL), plasmid-mediated AmpC (pAmpCs), and/or carbapenemase genes (5, 7, 12, 14, 18). Moreover, although it is known that the prevalence of intestinal colonization with ESC-R-Ent in returning travelers is very high (19), data regarding the *mcr-1*-Ent are needed. Only Arcilla et al. have explored this phenomenon, indicating that 0.9% of the ESBL producers isolated from stools of Dutch travelers co-possessed *mcr-1* (8).

We analyzed the pre- and posttrip stool samples (both within 1 week) of 38 people living in Switzerland and traveling to India during January to August 2015. A questionnaire was also filled out at each sampling time indicating that in the 12 months before going to India, participants frequently visited other countries but never suffered diarrhea. On the other hand, after the journey to India (mean stay, 17.8 days), 39% of the travelers suffered from travelers' diarrhea and additional symptoms, although antibiotics were not taken (see Table S1 in the supplemental material).

To detect ESC-R-Ent, ~20 μ g of fresh stools was enriched overnight in 10 ml LB broth containing a 10- μ g disk of cefuroxime. Then, 100 μ l was plated on BLSE, ChromID ESBL (bioMérieux), and Supercarba selective plates (20). After overnight incubation, colonies were identified using the matrix-assisted laser desorption/ionization–time of flight mass spectrometry (MALDI-TOF MS) (Bruker). Microdilution GN2F panels (Trek Diagnostics) were used to obtain the antibiotic MICs. The CT103XL microarray was used to detect *bla* genes. Multilocus sequence typing (MLST) for *E. coli* and *Klebsiella pneumoniae* (Warwick and Pasteur schemes, respectively) was used to deter-

mine the sequence type (ST); the *E. coli* phylogenetic group was also determined (21). The remaining stool samples were stored at -80°C without cryoprotectant.

Based on this initial screening, three (7.9%) people had stool samples positive for ESBL-producing *E. coli* before traveling, whereas 29 (76.3%) of them returned from India colonized with ESC-R-Ent, such as 26 CTX-M- and 3 DHA-producing *E. coli*. No carbapenemase producers were detected (Table 1). Overall, *E. coli* strains found in returning travelers were of non-hyperendemic STs and were mainly of phylogenetic groups A1 ($n = 11$) and B1 ($n = 8$). Notably, all of these ESC-R-Ent had polymyxin MICs of ≤ 0.5 $\mu\text{g/ml}$ and did not contain *mcr-1* as determined by PCR (1).

To improve our ability to detect possible Col-R-Ent (including those non-ESC-R), stored stools were enriched in 10 ml LB broth with two 10- μ g disks of colistin. After overnight incubation, 100 μ l was spread on four agar plates: MacConkey agar without or with colistin (4 $\mu\text{g/ml}$) and CHROMagar Orientation plus colistin (4 $\mu\text{g/ml}$) and vancomycin (8 $\mu\text{g/ml}$) without or with cefotaxime (2 $\mu\text{g/ml}$). MacConkey plates yielded numerous undistinguishable colonies of species naturally resistant to polymyxins (e.g., *Proteus* spp. and *Serratia marcescens*). In contrast, thanks to the typical color appearance of the different Gram-negative organisms, the CHROMagar Orientation plates allowed us to clearly identify Col-R *E. coli* (Col-R-Ec) colonies.

As depicted in Table 1 (rows highlighted in gray), only one

Received 15 April 2016 Returned for modification 19 May 2016

Accepted 28 May 2016

Accepted manuscript posted online 13 June 2016

Citation Bernasconi OJ, Kuenzli E, Pires J, Tinguely R, Carattoli A, Hatz C, Perreten V, Endimiani A. 2016. Travelers can import colistin-resistant *Enterobacteriaceae*, including those possessing the plasmid-mediated *mcr-1* gene. Antimicrob Agents Chemother 60:5080–5084. doi:10.1128/AAC.00731-16.

Address correspondence to Andrea Endimiani, aendimiani@gmail.com.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.00731-16>.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

TABLE 1 Extended-spectrum cephalosporin- and/or colistin-resistant *Enterobacteriaceae* detected in stool samples from 38 people before and after traveling to India during 2015^a

| Result from stool samples before traveling | | | | | | | | | | | | | Result from stool samples after traveling | | | | | | | | | | | | |
|--|---------|----------------------|---------------------------------|--------------------------|-----|-----|-----------------|-------------------------|-------|-------------------------|----------------------------|------|---|---|---|---|---|---|---|--|--|--|--|--|--|
| ID | Age/sex | No. of days in India | Species detected (strain) | bla gene(s) ^b | ST | CC | Phy | MIC, µg/ml ^c | | | PCR ⁺ for mcr-1 | | | | | | | | | | | | | | |
| | | | | | | | | COL | PoLB | MIC, µg/ml ^c | COL | PoLB | PCR ⁺ for mcr-1 | | | | | | | | | | | | |
| 3 ^d | 49/m | 31 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 8 ^e | 26/m | ND | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 9 | 25/m | 8 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 14 | 44/f | 15 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 18 | 52/f | 9 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 19 | 53/f | 20 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 23 ^h | 67/f | 14 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 26 ^e | 50/f | 14 | <i>E. coli</i> (26- <i>Ec</i>) | - | 69 | 69 | D1 | >4 (4) | >4 | - | - | - | - | - | - | - | - | - | - | | | | | | |
| 29 | 31/m | 15 | <i>E. coli</i> (29- <i>Ec</i>) | CTX-M-gr.9, TEMwt | 648 | 648 | D2 | ≤0.25 | ≤0.25 | - | - | - | - | - | - | - | - | - | - | | | | | | |
| 31 ^f | 40/m | 14 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 33 | 40/f | 32 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 34 | 36/f | 13 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 37 | 43/f | 27 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 41 | 68/f | 15 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 43 | 37/f | 17 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 45 ^f | 56/f | 14 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 50 | 33/m | 30 | <i>E. coli</i> (50- <i>Ec</i>) | CTX-M-15-like | 131 | 131 | B2 ₃ | ≤0.25 | ≤0.25 | - | - | - | - | - | - | - | - | - | - | | | | | | |
| 51 | 49/m | 19 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 52 ^g | 29/f | 14 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 55 ^d | 46/f | 31 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 56 | 42/m | 9 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 59 ^e | 50/f | 14 | <i>E. coli</i> (59- <i>Ec</i>) | CTX-M-15-like, TEMwt | 652 | NA | A1 | ≤0.25 | 0.5 | - | - | - | - | - | - | - | - | - | - | | | | | | |
| 60 | 53/f | 10 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 61 | 42/m | 9 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 63 | 33/f | 15 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 66 | 45/f | 18 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 68 | ND/m | 20 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 70 | 30/m | 27 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 71 | 59/f | 16 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 73 ^h | 53/f | 15 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 78 | 57/f | 20 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 80 | 26/f | 19 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 83 | 34/f | 23 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |
| 86 | 53/f | 15 | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | — | | | | | | |

| | | | | | | | | | | | |
|-----|------|----|---|---|--------------------------------------|------------------|-----|----|---------|-------|---|
| 88 | 25/m | 8 | — | <i>E. coli</i> (88R- <i>Ec</i> -ChromI) | CTX-M-15-like, TEMwt | 48 | 10 | A1 | ≤0.25 | ≤0.25 | — |
| | | | | <i>E. coli</i> (88R- <i>Ec</i>) | CTX-M-15-like | New ⁱ | NA | D1 | >4 (4) | >4 | — |
| 96 | 46/f | 35 | — | <i>E. coli</i> (96R- <i>Ec</i>) | CTX-M-15-like | 1491 | NA | A1 | ≤0.25 | ≤0.25 | — |
| 97 | 42/f | 20 | — | <i>E. coli</i> (97R- <i>Ec</i>) | CTX-M-15-like, TEMwt, TEM-11-like | 2914 | NA | D2 | ≤0.25 | 0.5 | — |
| | | | | <i>E. coli</i> (100R- <i>Ec</i> -Dri ^g) | CTX-M-gr.1, TEMwt | 155 | 155 | B1 | ≤0.25 | ≤0.25 | — |
| 100 | 57/f | 14 | — | <i>E. coli</i> (100R- <i>Ec</i>) | TEMwt | ST10 | 10 | A1 | >4 (12) | >4 | + |

^a Abbreviations used throughout: ID, identification code for the traveler; m, male; f, female; ND, not defined; ST, sequence type; CC, clonal complex; Phy, phylogenetic group; COL, colistin; PolB, polymyxin B; —, negative; +, positive; NA, not available or not applicable; wt, wild type (non-ESBL). Data highlighted in gray represent the colistin-resistant strains. All of these strains were detected with the CHROMagar Orientation plates containing colistin (4 µg/ml) plus vancomycin (8 µg/ml) with or without cefotaxime (2 µg/ml).

^b Based on the CheckPoints CT103X1 microarray results.

^c MICs were obtained in microdilution with the GN2F Trek panels; data in parentheses represent Etest results. For both methodologies, the MICs were obtained using colonies that grow on MacConkey agar plates without antibiotics.

^d Relatives or couples traveling together.

^e Relatives or couples traveling together.

^f Relatives or couples traveling together.

^g Friends traveling together.

^h Friends traveling together.

ⁱ The new STs could not be assigned because whole-genome sequencing for the strains was not performed as requested by the curator of the *E. coli* MLST scheme (see Table S3 in the supplemental material).

person was colonized with a Col-R-*Ec* strain before traveling, whereas four (10.5%) returning travelers had stools positive for Col-R-*Ec* (of which three coproduced ESBLs or pAmpCs). Remarkably, two (26-*Ec* and 100R-*Ec*) out of the overall five Col-R-*Ec* strains were non-ESC-R (see Table S2 in the supplemental material) and were identified only with the CHROMagar Orientation plates containing colistin but not cefotaxime. Notably, Col-R *Klebsiella* spp., *Enterobacter cloacae*, *Enterobacter aerogenes*, and *Salmonella* species strains were not found (2, 22). We cannot exclude that storing the stools at −80°C could influence the viability of Col-R strains.

Overall, one (2.6%) ST10/A1 *E. coli* strain (100R-*Ec*) was PCR positive for *mcr-1* (Table 1). This strain had a colistin MIC of 12 µg/ml by Etest and was coresistant to quinolones, trimethoprim-sulfamethoxazole, and tetracyclines (see Table S2 in the supplemental material). The strain carried HI2, FII, FIB, and X1 plasmid replicon types (PBRT kit; Diateva). Experiments with transformation into chemically competent DH5α *E. coli* failed to obtain *mcr-1*-carrying cells (23). Conjugation experiments using *E. coli* recipient strain JF33 (rifampin resistant) performed at 25 and 37°C and with selection on MacConkey plates containing colistin (2 µg/ml) and rifampin (50 µg/ml) were also unsuccessful.

To complete the bacterial typing of the *mcr-1*-positive strain and to identify the resistome and mobilome, raw genome data (MiSeq; Illumina) were screened by ResFinder-2.1 and PlasmidFinder-1.3 (24, 25). 100R-*Ec* carried resistance genes for aminoglycosides [*aadA1*, *aadA2*, and *aph(3')-Ia*], β-lactams [*bla*_{TEM-1}], polymyxins (*mcr-1*), quinolones (*qnrS1*), macrolides [*mph(A)*], phenicols (*cmlA1*), sulfonamides (*sul3*), tetracyclines [*tet(A)*], and trimethoprim (*dfrA12* and *dfrA14*).

By PlasmidFinder, the presence of IncF, IncX1, and IncHI2 plasmids was confirmed in the 100R-*Ec* strain. The *mcr-1* gene was identified at the end of a 26-kb contig (GenBank accession no. KX090925), flanked by two copies of IS*Apl1* in the opposite orientation forming a 4,211-bp IS*Apl1*-*mcr-1*-IS*Apl1* element. The 22,133-bp sequences preceding the IS*Apl1*-*mcr-1*-IS*Apl1* element shared >99% identity with DNA sequences of large IncHI2 plasmids, such as pKST313 from *Salmonella enterica* serotype Typhimurium (GenBank accession no. LN794248), pAPEC-O1-R from avian-pathogenic *E. coli* (GenBank accession no. DQ517526) (26, 27), and pSA26 carrying IS*Apl1*-*mcr-1* from an ST68 *E. coli* strain (GenBank accession no. KU743384). The *mcr-1* element was inserted into the *tra1* locus of the IncHI2 plasmid backbone immediately downstream of the conjugative protein gene *trhY*. The disruption of the *tra* region may explain the defective conjugation ability of this plasmid. Notably, the presence of *mcr-1* between duplicated IS*Apl1* insertion elements has so far only been observed in one human KPC-2-producing *E. coli* isolate in Germany and two IncHI2 plasmids isolated in the United Kingdom (4, 10).

Taking into account that 100R-*Ec* was of ST10/A1, and due to the high identity of the *mcr-1*-containing element with plasmids found in bacteria from animal origin, we speculate that this Col-R *E. coli* strain was acquired during the trip in India via the food chain or from environmental sources. Remarkably, the traveler was cocolonized with an ST155 CTX-M-producing *E. coli* strain (100R-*Ec*-Drig). A follow-up screening of the stool samples 3 and 6 months after returning from the trip indicated that the person was no longer colonized with both *E. coli* strains.

The remaining *mcr-1*-negative but Col-R *E. coli* strains (26-*Ec*, 78R-*Ec*, 86R-*Ec*, and 88R-*Ec*) underwent conjugation experi-

ments, but all failed to demonstrate a hypothetical plasmid-mediated mechanism of colistin resistance. We should note that such strains have been previously observed in humans and animals (8, 9, 28) and that they are Col-R, probably due to chromosomal mutations (10, 29). Notably, the stool samples from three of the travelers mentioned above were negative for Col-R *Ent* at the 3- and 6-month follow-up screenings. (ID-78 did not submit the stool samples for screening.) The risk factors determining the intestinal colonization with Col-R-*Ec* of people visiting India are unclear and should be investigated in the near future.

Our results indicate that the prevalence of intestinal *mcr-1-Ent* is probably underestimated because some isolates carrying only *mcr-1* (like 100R-*Ec*) could be routinely undetected when the screening is targeting the ESC-R and/or carbapenem-resistant strains (5, 7, 8, 12, 14, 18). It is possible that the IS*Apl1-mcr-1*-IS*Apl1* element present in 100R-*Ec* possesses the ability to transpose frequently, moving the *mcr-1* gene into the IncHI2 scaffold but also promoting its transposition in different plasmids. Multifocal plasmid types (IncP, IncX4, and IncI2) carrying *mcr-1*, but not *bla* genes conferring resistance to extended-spectrum cephalosporins (ESCs), have been previously reported, and some of them demonstrated a great potential for dissemination by conjugation among diverse Gram-negative species of the gut flora (1, 4, 18). Such colonizing bacteria may subsequently develop into difficult-to-treat infections, as observed for the ESBL producers acquired by people traveling from low- to high-prevalence countries (30–32).

The spread of non-ESC-R *mcr-1-Ent* could contribute to the silent expansion of this life-threatening resistance gene in both human and nonhuman settings. Specific and sensitive surveillance programs should be rapidly implemented to prevent unexpected outbreaks due to *mcr-1-Ent*.

ACKNOWLEDGMENTS

This work was supported by the Swiss National Science Foundation (SNF; grant no. 153377 to A.E.). The study was approved by the Ethikkommission Nordwest- und Zentralschweiz (EKNZ 239/12). Odette J. Bernasconi is a Ph.D. student (2015 to 2018) supported by the Hans Sigrist Foundation (Bern, Switzerland). João Pires is a Ph.D. student (2014 to 2017) supported by the SNF.

We thank Sara Kasraian and Alexandra Collaud for technical help.

FUNDING INFORMATION

This work, including the efforts of Andrea Endimiani, was funded by Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (SNF) (153377).

REFERENCES

- Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J. 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* 16:161–168. [http://dx.doi.org/10.1016/S1473-3099\(15\)00424-7](http://dx.doi.org/10.1016/S1473-3099(15)00424-7).
- Skov RL, Monnet DL. 2016. Plasmid-mediated colistin resistance (*mcr-1* gene): three months later, the story unfolds. *Euro Surveill* <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.9.30155>.
- Poirel L, Kieffer N, Liassine N, Thanh D, Nordmann P. 2016. Plasmid-mediated carbapenem and colistin resistance in a clinical isolate of *Escherichia coli*. *Lancet Infect Dis* 16:281. [http://dx.doi.org/10.1016/S1473-3099\(16\)00006-2](http://dx.doi.org/10.1016/S1473-3099(16)00006-2).
- Falgenhauer L, Waezsada SE, Yao Y, Imirzalioglu C, Kasbohrer A, Roesler U, Michael GB, Schwarz S, Werner G, Kreienbrock L, Chakraborty T. 2016. Colistin resistance gene *mcr-1* in extended-spectrum β -lactamase-producing and carbapenemase-producing Gram-negative bacteria in Germany. *Lancet Infect Dis* 16:282–283. [http://dx.doi.org/10.1016/S1473-3099\(16\)00009-8](http://dx.doi.org/10.1016/S1473-3099(16)00009-8).
- Du H, Chen L, Tang YW, Kreiswirth BN. 2016. Emergence of the *mcr-1* colistin resistance gene in carbapenem-resistant *Enterobacteriaceae*. *Lancet Infect Dis* 16:287–288. [http://dx.doi.org/10.1016/S1473-3099\(16\)00056-6](http://dx.doi.org/10.1016/S1473-3099(16)00056-6).
- Olaitan AO, Chabou S, Okdah L, Morand S, Rolain JM. 2016. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis* 16:147. [http://dx.doi.org/10.1016/S1473-3099\(15\)00540-X](http://dx.doi.org/10.1016/S1473-3099(15)00540-X).
- Hasman H, Hammerum AM, Hansen F, Hendriksen RS, Olesen B, Agerso Y, Zankari E, Leekitcharoenphon P, Stegger M, Kaas RS, Cavaco LM, Hansen DS, Aarestrup FM, Skov RL. 2015. Detection of *mcr-1* encoding plasmid-mediated colistin-resistant *Escherichia coli* isolates from human bloodstream infection and imported chicken meat, Denmark 2015. *Euro Surveill* <http://dx.doi.org/10.2807/1560-7917.ES.2015.20.49.30085>.
- Arcilla MS, van Hattem JM, Matamoros S, Melles DC, Penders J, de Jong MD, Schultz C. 2016. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis* 16:147–149. [http://dx.doi.org/10.1016/S1473-3099\(15\)00541-1](http://dx.doi.org/10.1016/S1473-3099(15)00541-1).
- Cannatelli A, Giani T, Antonelli A, Principe L, Luzzaro F, Rossolini GM. 2016. First detection of the *mcr-1* colistin resistance gene in *Escherichia coli*, Italy. *Antimicrob Agents Chemother* 60:3257–3258. <http://dx.doi.org/10.1128/AAC.00246-16>.
- Doumith M, Godbole G, Ashton P, Larkin L, Dallman T, Day M, Muller-Pebody B, Ellington MJ, de Pinna E, Johnson AP, Hopkins KL, Woodford N. 18 April 2016. Detection of the plasmid-mediated *mcr-1* gene conferring colistin resistance in human and food isolates of *Salmonella enterica* and *Escherichia coli* in England and Wales. *J Antimicrob Chemother* <http://dx.doi.org/10.1093/jac/dkw093>.
- Suzuki S, Ohnishi M, Kawanishi M, Akiba M, Kuroda M. 2016. Investigation of a plasmid genome database for colistin-resistance gene *mcr-1*. *Lancet Infect Dis* 16:284–285. [http://dx.doi.org/10.1016/S1473-3099\(16\)00008-6](http://dx.doi.org/10.1016/S1473-3099(16)00008-6).
- Malhotra-Kumar S, Xavier BB, Das AJ, Lammens C, Hoang HT, Pham NT, Goossens H. 2016. Colistin-resistant *Escherichia coli* harbouring *mcr-1* isolated from food animals in Hanoi, Vietnam. *Lancet Infect Dis* 16:286–287. [http://dx.doi.org/10.1016/S1473-3099\(16\)00014-1](http://dx.doi.org/10.1016/S1473-3099(16)00014-1).
- Haenni M, Poirel L, Kieffer N, Chatre P, Saras E, Metayer V, Dumoulin R, Nordmann P, Madec JY. 2016. Co-occurrence of extended spectrum β lactamase and MCR-1 encoding genes on plasmids. *Lancet Infect Dis* 16:281–282. [http://dx.doi.org/10.1016/S1473-3099\(16\)00007-4](http://dx.doi.org/10.1016/S1473-3099(16)00007-4).
- Perrin-Guyomard A, Bruneau M, Houe P, Deleurme K, Legrandois P, Poirier C, Soumet C, Sanders P. 2016. Prevalence of *mcr-1* in commensal *Escherichia coli* from French livestock, 2007 to 2014. *Euro Surveill* <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.6.30135>.
- Zurfuh K, Poirel L, Nordmann P, Nuesch-Inderbinen M, Hachler H, Stephan R. 2016. Occurrence of the plasmid-borne *mcr-1* colistin resistance gene in ESBL-producing *Enterobacteriaceae* in river water and imported vegetable samples in Switzerland. *Antimicrob Agents Chemother* 60:2594–2595. <http://dx.doi.org/10.1128/AAC.00066-16>.
- Webb HE, Granier SA, Marault M, Millemann Y, den Bakker HC, Nightingale KK, Bugarel M, Ison SA, Scott HM, Loneragan GH. 2016. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis* 16:144–145. [http://dx.doi.org/10.1016/S1473-3099\(15\)00538-1](http://dx.doi.org/10.1016/S1473-3099(15)00538-1).
- Kluytmans-van den Bergh MF, Huizinga P, Bonten MJ, Bos M, De Bruyne K, Friedrich AW, Rossen JW, Savelkoul PH, Kluytmans JA. 2016. Presence of *mcr-1*-positive *Enterobacteriaceae* in retail chicken meat but not in humans in the Netherlands since 2009. *Euro Surveill* <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.9.30149>.
- Yao X, Doi Y, Zeng L, Lv L, Liu JH. 2016. Carbapenem-resistant and colistin-resistant *Escherichia coli* co-producing NDM-9 and MCR-1. *Lancet Infect Dis* 16:288–289. [http://dx.doi.org/10.1016/S1473-3099\(16\)00057-8](http://dx.doi.org/10.1016/S1473-3099(16)00057-8).
- Kuenzli E, Jaeger VK, Frei R, Neumayr A, DeCrom S, Haller S, Blum J, Widmer AF, Furrer H, Battegay M, Endimiani A, Hatz C. 2014. High colonization rates of extended-spectrum β -lactamase (ESBL)-producing *Escherichia coli* in Swiss travellers to South Asia—a prospective observational multicentre cohort study looking at epidemiology, microbiology and risk factors. *BMC Infect Dis* 14:528. <http://dx.doi.org/10.1186/1471-2334-14-528>.
- Viau R, Frank KM, Jacobs MR, Wilson B, Kaye K, Donskey CJ, Perez

- F, Endimiani A, Bonomo RA. 2016. Intestinal carriage of carbapenemase-producing organisms: current status of surveillance methods. *Clin Microbiol Rev* 29:1–27. <http://dx.doi.org/10.1128/CMR.00108-14>.
21. Escobar-Paramo P, Le Menac'h A, Le Gall T, Amorin C, Gouriou S, Picard B, Skurnik D, Denamur E. 2006. Identification of forces shaping the commensal *Escherichia coli* genetic structure by comparing animal and human isolates. *Environ Microbiol* 8:1975–1984. <http://dx.doi.org/10.1111/j.1462-2920.2006.01077.x>.
 22. Zeng KJ, Doi Y, Patil S, Huang X, Tian GB. 2016. Emergence of plasmid-mediated *mcr-1* gene in colistin-resistant *Enterobacter aerogenes* and *Enterobacter cloacae*. *Antimicrob Agents Chemother* 60:3862–3863. <http://dx.doi.org/10.1128/AAC.00345-16>.
 23. Carattoli A, Seiffert SN, Schwendener S, Perreten V, Endimiani A. 2015. Differentiation of IncL and IncM plasmids associated with the spread of clinically relevant antimicrobial resistance. *PLoS One* 10:e0123063. <http://dx.doi.org/10.1371/journal.pone.0123063>.
 24. Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, Moller Aarestrup F, Hasman H. 2014. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <http://dx.doi.org/10.1128/AAC.02412-14>.
 25. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <http://dx.doi.org/10.1093/jac/dks261>.
 26. Kariuki S, Okoro C, Kiiru J, Njoroge S, Omuse G, Langridge G, Kingsley RA, Dougan G, Revathi G. 2015. Ceftriaxone-resistant *Salmonella enterica* serotype Typhimurium sequence type 313 from Kenyan patients is associated with the *bla*_{CTX-M-15} gene on a novel IncHI2 plasmid. *Antimicrob Agents Chemother* 59:3133–3139. <http://dx.doi.org/10.1128/AAC.00078-15>.
 27. Johnson TJ, Wannemuehler YM, Scaccianoce JA, Johnson SJ, Nolan LK. 2006. Complete DNA sequence, comparative genomics, and prevalence of an IncHI2 plasmid occurring among extraintestinal pathogenic *Escherichia coli* isolates. *Antimicrob Agents Chemother* 50:3929–3933. <http://dx.doi.org/10.1128/AAC.00569-06>.
 28. Malhotra-Kumar S, Xavier BB, Das AJ, Lammens C, Butaye P, Goossens H. 2016. Colistin resistance gene *mcr-1* harboured on a multidrug resistant plasmid. *Lancet Infect Dis* 16:283–284. [http://dx.doi.org/10.1016/S1473-3099\(16\)00012-8](http://dx.doi.org/10.1016/S1473-3099(16)00012-8).
 29. Quesada A, Porrero MC, Tellez S, Palomo G, Garcia M, Dominguez L. 2015. Polymorphism of genes encoding PmrAB in colistin-resistant strains of *Escherichia coli* and *Salmonella enterica* isolated from poultry and swine. *J Antimicrob Chemother* 70:71–74. <http://dx.doi.org/10.1093/jac/dku320>.
 30. Laupland KB, Church DL, Vidakovich J, Mucenski M, Pitout JD. 2008. Community-onset extended-spectrum β -lactamase (ESBL) producing *Escherichia coli*: importance of international travel. *J Infect* 57:441–448. <http://dx.doi.org/10.1016/j.jinf.2008.09.034>.
 31. Banerjee R, Strahilevitz J, Johnson JR, Nagwekar PP, Schora DM, Shevrin I, Du H, Peterson LR, Robicsek A. 2013. Predictors and molecular epidemiology of community-onset extended-spectrum beta-lactamase-producing *Escherichia coli* infection in a Midwestern community. *Infect Control Hosp Epidemiol* 34:947–953. <http://dx.doi.org/10.1086/671725>.
 32. Soraas A, Sundsfjord A, Sandven I, Brunborg C, Jenum PA. 2013. Risk factors for community-acquired urinary tract infections caused by ESBL-producing enterobacteriaceae—a case-control study in a low prevalence country. *PLoS One* 8:e69581. <http://dx.doi.org/10.1371/journal.pone.0069581>.